

# NOTICE OF PUBLICATION



AQUAFISH COLLABORATIVE RESEARCH SUPPORT PROGRAM

## RESEARCH REPORTS

Sustainable Aquaculture for a Secure Future

**Title:** Growth performance, survival, feed utilization and nutrient utilization of African catfish (*Clarias gariepinus*) larvae co-fed *Artemia* and a micro-diet containing freshwater atyid shrimp (*Caridina nilotica*) during weaning

**Author(s):** Chepkirui-Boit, V.<sup>1</sup>, Ngugi, C.C.<sup>1</sup>, Bowman, J.<sup>2</sup>, Oyoo-Okoth, E.<sup>1,3</sup>, Rasowo, J.<sup>4</sup>, Mugo-Bundi, J.<sup>1</sup> and Cherop, L.<sup>1</sup>

<sup>1</sup>Department of Fisheries and Aquatic Sciences, Moi University, Eldoret, Kenya

<sup>2</sup>Department of Wildlife, Oregon State University, Corvallis, OR, USA

<sup>3</sup>Department of Aquatic Ecology and Ecotoxicology, Amsterdam, The Netherlands

<sup>4</sup>Department of Biological Sciences, Moi University, Eldoret, Kenya

**Date:** September 3, 2010

Publication Number: CRSP Research Report 10-258

The CRSP will not be distributing this publication. Copies may be obtained by writing to the authors.

**Abstract:** Problems of limited number of dry feeds as supplement or replacement of live feeds have led to poor larval nutrition in many species of fish. Therefore, the suitability of co-feeding 8-day-old African catfish (*Clarias gariepinus*) posthatch larvae using live feed (*Artemia salina*) and formulated dry diet containing freshwater atyid shrimp (*Caridina nilotica*) during weaning was investigated. The experiment ended after 21 days of culture and respective groups compared on the basis of growth performance, survival, feed utilization and nutrient utilization. Larvae co-fed using 50% *Artemia* and 50% formulated dry diet resulted in significantly ( $P < 0.05$ ) better growth performance, food gain ratio (FGR), protein efficiency ratio (PER) and productive protein values (PPV) than other treatments. The lowest growth performance occurred in larvae weaned using 100% formulated and commercial dry diets. Better survival of over 90% was obtained in larvae weaned using 50% *Artemia* and 50% dry diet, while abrupt weaning using 100% dry diets resulted in lower survival (<75%). These results support a recommendation of co-feeding *C. gariepinus* larvae using a formulated dry diet containing *C. nilotica* and 50% live feed when weaning is performed after 8 days posthatching period.

This abstract was excerpted from the original paper, which was published in African Journal of Microbiology Research, Aquaculture Nutrition 2010: 1-8.

**CRSP RESEARCH REPORTS** are published as occasional papers by the Management Entity, AquaFish Collaborative Research Support Program, Oregon State University, 418 Snell Hall, Corvallis, Oregon 97331-1643 USA. The AquaFish CRSP is supported by the US Agency for International Development under CRSP Grant No. EPP-A-00-06-00012-00. See the website at <aquafishcrsp.oregonstate.edu>.