

NOTICE OF PUBLICATION



AQUAFISH COLLABORATIVE RESEARCH SUPPORT PROGRAM

RESEARCH REPORTS

Sustainable Aquaculture for a Secure Future

Title: Effects of addition of red tilapia (*Oreochromis* spp.) at different densities and sizes on production, water quality and nutrient recovery of intensive culture of white shrimp (*Litopenaeus vannamei*) in cement tanks

Author(s): Yuan, D.^a, Yi, Y.^b, Yakupitiyage, A.^c, Fitzimmons, K.^d and Diana, J.^e
^a Network of Aquaculture Centres in Asia-Pacific, Bangkok 10903, Thailand
^b Division of Aquaculture, E-Institute, Shanghai Ocean University, Shanghai 200090, China
^c Aquaculture and Aquatic Resources Management, School of Environment, Resources and Development, Asian Institute of Technology, Pathumthani 12120, Thailand
^d Environmental Research Lab, Department of Soil, Water and Environmental Science, University of Arizona, Tucson, USA
^e School of Natural Resources and Environment, University of Michigan, Ann Arbor, Michigan, USA

Date: June 4, 2010 Publication Number: CRSP Research Report 10-257

The CRSP will not be distributing this publication. Copies may be obtained by writing to the authors.

Abstract: An experiment was conducted in 21 outdoor cement tanks (2.5×2×1.2 m) from 8 December 2005 to 3 March 2006 to determine the effects of adding red tilapia (*Oreochromis* spp.) at different densities and sizes on production, water quality and nutrient recovery in intensive culture tanks of white shrimp (*Litopenaeus vannamei*). Shrimp postlarvae of 0.06 g were stocked into all tanks at a density of 60 postlarvae m⁻², while either small (13.8±0.2g) or large (41.9 ± 0.3 g) mono-sex tilapia fingerlings were stocked into the shrimp tanks two weeks later at low (0.4fish m⁻²), medium (0.8 fish m⁻²) or high (1.2 fish m⁻²) density. Water depth in all tanks was maintained at 1 m and salinity at 20 ppt. Water loss due to evaporation was compensated weekly. The experiment was conducted in a 2 × 3 factorial design, while three additional tanks for shrimp monoculture were set as a control. All treatments and the control were randomly allocated to tanks in triplicate each. Shrimps were fed three times daily with commercial pellets using feeding trays made with metal frame and nylon mesh (0.6× 0.6× 0.05 m) at the same feeding rates as those for the control. No separate feed was given to tilapia.
The highest shrimp survival rate of 66.8% was obtained in the small-low density tilapia treatment, which was significantly higher than those in other treatments and the control. The small-low density tilapia treatment had the highest shrimp yield and lowest feed conversion ratio, which was similar to those in the control and the large-low and small-

CRSP RESEARCH REPORTS are published as occasional papers by the Management Entity, AquaFish Collaborative Research Support Program, Oregon State University, 418 Snell Hall, Corvallis, Oregon 97331-1643 USA. The Aquafish CRSP is supported by the US Agency for International Development under CRSP Grant No. EPP-A-00-06-00012-00. See the website at <aquafishcrsp.oregonstate.edu>.

Continued...

medium density tilapia treatments, but significantly better than those in other treatments. Factorial analyses revealed that the increase of tilapia density from 0.4 to 1.2 fish m⁻² and size from 13.8 to 41.9 g negatively affected shrimp production performance but remarkably increased the combined production of shrimp and tilapia. Polyculture incorporated 36.0-49.5% of the total nitrogen input and 14.2-26.5% of the total phosphorous input into shrimp and tilapia, which were significantly higher than those (27.1% and 8.9%) in the monoculture, respectively. The nutrient recovery efficiency increased with increased tilapia stocking size and density. Polyculture with small tilapia stocked at low density had the best economic performance among all treatments and control, and significantly better than small-high, large-medium and large-high density tilapia treatments.

It was concluded that addition of red tilapia at suitable stocking densities and sizes into intensive white shrimp monoculture can improve productivity, profitability, nutrient utilization and environmental friendliness of shrimp monoculture. The suitable stocking density and size of red tilapia identified in this study were 0.4 fish m⁻² and 13.7 g respectively. Red tilapia could be stocked at higher density and larger size up to 1.2 fish m⁻² and 42 g respectively to maximize system productivity and minimize nutrient waste without affecting shrimp survival, but economic performance could be negatively affected. Shrimp-tilapia polyculture should be promoted to improve sustainability of shrimp culture.

This abstract was excerpted from the original paper, which was published in *Aquaculture* 298: 226–238.